

West Michigan Veterinary Service
DAIRY VACCINE/HEALTH RECOMMENDATIONS

BIRTH

1. Colostrum (6qt in 6 hr.)
2. "Bo-Se" IM (3cc)
3. Coat Navel in strong iodine (3days)
4. Vitamin A&D IM (2cc)
5. "Calf-Guard" orally 30 minutes before colostrum if not using Scourguard 4KC

1 to 3 DAYS

1. Inforce-3 (IN) intranasal

UP TO 4 WEEKS

1. Dehorn. (Horn Stop, burner, paste)
2. Remove extra teats.
3. Check navel for drainage or swelling.

3-6 WEEKS

1. Reinject Viramin A&D and "Bo-Se" 3cc IM (optional).
2. "Pulmoguard PHM" or "One Shot" 2cc SubQ.
3. "Vision 7w/somnus" 2cc Sub-Q.

3 MONTHS

1. Bovishield Gold 5+L5 FP+/ Spirovac 2cc IM. 3. "One Shot" or PMG-PHM
2. "Vision 7 w/somnus" 2ccSub-Q. 4. Dehorn. (if not done earlier)

4 MONTHS

1. Bovishield Gold 5+L5 FP+/ Spirovac 2cc IM
2. "Vision 7 w/somnus" 2ccSub-Q.
3. "One Shot" or PMG-PHM
4. Deworm.

12-14 MONTHS

1. Bovishield Gold 5+L5 FP+/ Spirovac 2cc IM
2. "Vision 7 w/somnus" 2ccSub-Q. 4. Deworm.
3. "One Shot" or PMG-PHM

3 MONTHS PRIOR TO CALVING

1. "J-5 E. coli Mastitis vaccine. 5cc Sub-Q (optional)
2. "Scourguard 4KC" 2cc SubQ.

2 MONTHS PRIOR TO CALVING

1. "Bovine Pilishield C" or "Scourguard 4KC"
2. "J-5 E. coli Mastitis vaccine. 5cc Sub-Q (optional)
3. "Vision 7 w/somnus" 2ccSub-Q. (or once a year to all cows).

2-3 WEEKS PRIOR TO CALVING

1. "Mu-Se" IM or "Multimin" Sub-Q.
2. Deworm.

3-7 DAYS AFTER CALVING

1. "J-5 E. coli Mastitis vaccine. 5cc Sub-Q (optional)

2-5 WEEKS POST CALVING

1. Bovishield Gold 5+L5 FP+/ Spirovac 2cc IM. Or "CattleMaster 4+L5" and Spirovac once a year.

1 MONTH PRIOR TO DRY OFF

1. "J-5" optional or one month after dry off.
2. "Scourguard 4KC"

DRY OFF

1. Dry Treat Udder.
2. "Vision 7 w/somnus" or once a year.
3. "Orbeseal" teat sealant.
4. "Bovine Pilishield C" or "Scourguard 4KC"
5. "J-5" (optional)
6. Trim Feet.

2-3 WEEKS PRIOR TO CALVING

1. "Mu-Se" IM or "Multimin" Sub-Q.
2. Deworm. Or at dry off.

ONCE A YEAR

1. "One Shot" pneumonia vaccine. Whole herd (down to 6 months) in Fall.
2. "Vision 7 w/somnus" 2ccSub-Q. (or at dry off).
3. "CattleMaster 5+L5" if Bovishield Gold 5+L5 FP+/ Spirovac 2cc IM is not given to fresh cows.

TWICE A YEAR

1. "Leptoferm" (or booster when confirmed pregnant).

.....

- Salmonella (SRP), Pinkeye (SolidBac) vaccine and Autogenous vaccines are all optional. They should be given twice, a month apart. The best time to do these may be between 5 and 12 months of age.

- Pinkeye (SolidBac) does not need be boosted.

- Salmonella and Autogenous vaccines can be boosted at dry off.

- "Spirovac" herds need to add this at 3 and 4 months of age, with the Bovishield Gold 5+L5 FP+/ Spirovac 2cc IM. Also at 12-14 months and 2-5 weeks post calving. The combination vaccine is cheaper than giving both separately.

***However, please avoid giving over 3 vaccinations at one time.